

HARMFUL ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

zelam

TRIBUTE™ SC

SYSTEMIC FUNGICIDE

Controls many important foliar diseases of barley, oats and wheat, white rot in onions, powdery mildew in peas, and crown and stem rust in ryegrass seed crops

F U N G I C I D E

NET CONTENTS: 5 LITRES

TM Trademark of Tapuae Partnership NZ

Registered pursuant to the ACVM Act 1997 No. P7174 See www.nzfsa.govt.nz/acvm for registration conditions

ACTIVE INGREDIENT: Contains 375g/litre triadimenol in the form of a suspension concentrate.

DANGER: May cause organ damage from repeated oral exposure at high doses.

WARNING: May be harmful if swallowed, inhaled or absorbed through the skin.

PRECAUTIONS: Avoid contact with skin or inhalation of spray mist. Wear eye protection when handling the concentrate. Gloves should be worn at all times. Remove protective clothing and wash hands and face thoroughly before meals and after work. DO NOT eat, drink, or smoke while using.

FIRST AID: If swallowed, DO NOT induce vomiting.

Give nothing by mouth. Take to hospital without delay.

If skin contact occurs, remove contaminated clothing and wash affected areas of skin. DO NOT scrub the skin. Get medical attention if irritation persists. If splashed in eyes, hold lids open and flush with water for 15 minutes. Take to hospital immediately. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

ECOTOXIC: Toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment and terrestrial vertebrates.

DISPOSAL: Triple rinse container and add residue to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill. Avoid contamination of any waterway with product or empty container.

STORAGE: Store in original container, tightly closed, away from foodstuffs. Keep out of reach of children.

SHELF LIFE: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

REGISTERED TO: Tapuae Partnership,

Hudson Rd, New Plymouth, NZ
Tel: 06-755-9234 Fax: 06-755-1174

**EMERGENCY PHONE:
0800 CHEMCALL - 0800 243 6225**

**PLEASE READ ENTIRE LABEL BEFORE USE.
SHAKE OR STIR WELL BEFORE USE.**

| DISEASE | | DIRECTIONS FOR USE |
|---------------------|--|---|
| BARLEY | Scald, leaf rust | Use 330ml/ha and apply at first appearance of disease. A further application may be required 3 - 4 weeks later if disease pressure persists. |
| OATS | Crown rust | |
| RYEGRASS SEED CROPS | Crown rust, stem rust | |
| WHEAT | Stripe rust, leaf rust, powdery mildew, speckled leaf blotch | |
| ONIONS | White rot | Use 1 litre/ha. Apply 6 - 8 weeks after planting, then 2 further applications at 3 - 4 weekly intervals. The use of a suitable seed treatment is recommended. |
| PEAS | Powdery mildew | Use 165ml/ha and apply at first appearance of disease. Repeat 14 days later if disease persists. |

WITHHOLDING PERIOD: Peas - 28 days; Ryegrass seed crops and onions - 35 days; cereals - 49 days.

APPLICATION: For cereals, ryegrass and peas apply in at least 200 litres of water/ha by ground application or 50 litres/ha by aerial application. Use higher water volumes in dense or advanced crops especially under dry weather conditions. For onions apply 1000 litres of water/ha.

COMPATIBILITY: Compatible with most commonly used herbicides and insecticides.

MODE OF ACTION: Tribute™ SC has both contact and systemic action. Absorbed by leaves and stems and translocated in an upward and outward movement in the plant to provide some protection to new growth.

RESISTANCE: Resistance to this fungicide and related DMI products could develop in some disease situations from repeated use. Use a maximum of three applications per season. Resistance is most effectively prevented by adhering to full application rates and by mixing this product with an effective dose of a fungicide from a different cross-resistance group.

RAINFALL: Rain falling within 1 hour following treatment may reduce effectiveness.

MIXING: Pour the correct amount of Tribute™ SC into the partly filled spray tank and while agitating, add the remainder of the water.

NOTICE TO BUYER: The manufacturers and vendors of Tribute™ SC warrant the formulation to contain the specified active ingredient within accepted tolerance when packed but make no warranty of any kind on the use of this material and accept no responsibility for results, whether the material has been used in accordance with directions or not.

BATCH NO:

D.O.M:

