

HARMFUL
KEEP OUT OF REACH OF CHILDREN



TURFWET*

Soil Active Wetting Agent for Turf

**Enhances soil & thatch penetration.
Improves surface removal of water soil & plant
wetting aid. Improves spray coverage**

A D D I T I V E

NET CONTENTS: 5 LITRE

* Tradename of Tapuae Partnership, NZ.

PLEASE READ ENTIRE LABEL BEFORE MIXING OR USE

ACTIVE INGREDIENT: Contains 1000g/litre blended alkyl aryl polyoxyethylene glycols

WARNING: May be harmful if swallowed, inhaled or absorbed through the skin.

FIRST AID: If medical advice is needed : Have product container or label present. If swallowed, DO NOT induce vomiting. Give nothing by mouth. Take to hospital without delay. If skin contact occurs, remove contaminated clothing and wash affected areas of skin. DO NOT scrub the skin. Get medical attention if irritation persists. If concentrate is splashed in eyes, hold lids open and flush with water for 15 minutes. Take to hospital immediately. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

PERSONAL PROTECTION: Avoid contact with skin or inhalation of spray mist. Gloves should be worn at all times. Wear eye protection when handling the concentrate. Ensure adequate ventilation when handling the concentrate. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work. Do not eat, drink or smoke while using this product.

ECOTOXIC: Harmful to terrestrial vertebrates.

DISPOSAL: When empty triple rinse container and add residue to spray tank. Dispose of this product only by using according to the label, or at an approved landfill, or other approved facility. Avoid contamination of any water supply or stream with product or empty container.

STORAGE: Store in original container out of reach of children.

SHELF LIFE: When stored appropriately this product should show no significant degradation for 2 years from date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the MSDS available from Zelum Limited or www.zelum.co.nz

ERMA NEW ZEALAND: Additional information is available from the ERMA NZ website - www.ermanz.govt.nz. Approval Code: HSR002503

NOTICE TO BUYER: The manufacturer and vendors of **Turfwet** warrant the formulation to contain the specified active ingredient within accepted tolerance but make no warranty of any kind on the uses of material and accept no responsibility, whether the material is used in accordance with the label directions or not.

SITUATION	RATE	DIRECTIONS
Turf, Lawns, Cotula Greens	10 litres Turfwet in not less than 250 litres of water/ ha.	Apply as an initial application in August-September. Apply monthly repeat treatment of 5 litres per hectare as required.
Bowling greens	1.25 - 3 litres Turfwet per green	Apply once yearly. Repeat if necessary.
With Insecticides and fungicides	25ml of Turfwet per 100 litres of spray mix	Non-selective dessicant herbicides and grass killers may benefit from a use rate of 100-200 ml of Turfwet * per 100 litres of spray mix. Rainfall soon after treatment will not reduce effectiveness.
With Herbicides	50 - 100 ml of Turfwet per 100 litres of spray mix.	
Pot Plants	1 ml Turfwet per litre	Apply by soaking potted plant in the use solution. For large volumes, use a watering can to wet soil completely.

Mixing: Always add **Turfwet** to a near filled spray tank after the other products have been added. Ensure **Turfwet** is well mixed into the spray solution. Avoid foaming by adding remaining water below the surface. **Turfwet** should be applied in sufficient water to achieve uniform coverage. Ideally application should follow a light syringe of the turf.
Foaming: If foaming is unavoidable the use of Foamex*, an anti-foam agent is recommended.

GENERAL: **Turfwet** can be mixed with all commonly used fungicides, such as Taratek*5F, insecticides, and herbicides. It mixes readily with water. Under hardened growth conditions use the high rate of **Turfwet**. Use sufficient **Turfwet** to form a continuous film and give uniform coverage. Excessive rates may lead to undesirable foaming and unnecessary runoff. Because **Turfwet** can release some of a plants natural leaf waxes **Turfwet** should be applied in the late afternoon when temperatures are cooler to ensure no burning to sensitive leaves.

REGISTERED TO: Tapuae Partnership,
Hudson Road, New Plymouth, NZ.
Tel: 06 755-9234 Fax: 06 755-1174

EMERGENCY PHONE:
0800 CHEMCALL - 0800 243 6225



BATCH NO:

D.O.M: