

HARMFUL, ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

TRIBUTE[®]SC

FUNGICIDE

Contains: 375 g/litre triadimenol in the form of a suspension concentrate

**For disease control in barley, oats, wheat,
onions, peas and ryegrass seed crops**

®Registered Trademark of Zelum Limited



PLEASE READ THE ENTIRE LABEL BEFORE MIXING OR USE. SHAKE OR STIR WELL BEFORE USE

HAZARD CLASSIFICATION: 6.1D, 6.9A, 9.1B, 9.2C, 9.3C

WARNING: This product may be harmful if swallowed, inhaled or absorbed through the skin. May cause organ damage from repeated oral exposure at high doses.

FIRST AID: If swallowed, DO NOT induce vomiting. Give nothing by mouth. Take to hospital without delay. If skin contact occurs, remove contaminated clothing and wash affected areas of skin. DO NOT scrub the skin. Get medical attention if irritation persists. If splashed in eyes, hold lids open and flush with water for 15 minutes. Take to hospital immediately.

For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

PERSONAL PROTECTION: Avoid skin and eye contact and inhalation of spray mist. When mixing or applying wear waterproof gloves, eye protection, boots and overalls. Do not eat, drink or smoke while using. Wash splashes of concentrate from skin immediately. Ensure adequate ventilation when handling the concentrate. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work.

ECOTOXIC: Toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment and terrestrial vertebrates.

STORAGE: Store in original container, tightly closed and in a dry, cool area away from foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals.

DISPOSAL: Triple rinse empty container and add rinsate to the spray tank. If recycling, discard cap and deliver clean container to an Agrecovery depot. Alternatively crush and bury in an approved landfill.

Dispose of product only by using according to this label, or at an approved disposal facility.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the MSDS available from Zelam Ltd or www.zelam.com.

Approved pursuant to the HSNO Act 1996, No. HSR001615. See www.epa.govt.nz for approval conditions.

Registered pursuant to the ACVM Act 1997, No. P007174 See www.foodsafety.govt.nz for registration conditions.

It is an offence to use this product on animals,

DIRECTIONS FOR USE:

CROP	DISEASE	CRITICAL COMMENTS
BARLEY	Scald, leaf rust	Use 330 ml/ha and apply at first appearance of disease. A further application may be required 3 - 4 weeks later if disease pressure persists
OATS	Crown rust	
RYEGRASS SEED CROPS	Crown rust, stem rust	
WHEAT	Stripe rust, leaf rust, powdery mildew, speckled leaf blotch	
ONIONS	White rot	Use 1 litre/ha. Apply 6 - 8 weeks after planting, then 2 further applications at 3 - 4 weekly intervals. The use of a suitable seed treatment is recommended.
PEAS	Powdery mildew	Use 165 ml/ha and apply at first appearance of disease. Repeat 14 days later if disease persists.

APPLICATION: For cereals, ryegrass and peas apply in at least 200 litres of water/ha by ground application or 50 litres/ha by aerial application. Use higher water volumes in dense or advanced crops especially under dry weather conditions. For onions apply 1000 litres of water/ha.

COMPATIBILITY: Compatible with most commonly used herbicides and insecticides.

MODE OF ACTION: Tribute SC has both contact and systemic action. Absorbed by leaves and stems and translocated in an upward and outward movement in the plant to provide some protection to new growth.

MIXING: Pour the correct amount of Tribute SC into the partly filled spray tank and while agitating, add the remainder of the water.

Use an appropriate water rate for the weeds present but use a minimum of 200 litres/ha.

RAINFALL: Rain falling within 1 hour following treatment may reduce effectiveness.

WITHHOLDING PERIODS: Peas - 28 days; Ryegrass seed crops and onions - 35 days; cereals - 49 days.

It is an offence for users of this product to cause residues exceeding the relevant MRL in Food Notice: Maximum Residue Levels for Agricultural Compounds.

RESISTANCE MANAGEMENT: **GROUP 3 FUNGICIDE** Tribute contains a triazole fungicide with the demethylation inhibition (DMI) mode of action. Naturally occurring strains of disease causing fungi resistant to Tribute could develop. These resistant strains may dominate when Tribute and other related group 3 fungicides are used excessively. To minimise this risk use strictly in accordance with label instructions and resistance management strategies that may exist for particular diseases on this label. For further information see www.resistance.nzpps.org

NOTICE TO BUYER: The manufacturer and vendors of Tribute SC warrant the formulation to contain the specified active ingredient within accepted tolerance but make no warranty of any kind on the uses of material and accept no responsibility, whether the material is used in accordance with the label directions or not.